

Editorial: Changes in Forest Ecosystem Nutrition

Lang, Friederike; Kruger, Jaane; Kaiser, Klaus; Bol, Roland; Loeppmann, Sebastian

Frontiers in Forests and Global Change

DOI:

[10.3389/ffgc.2021.752063](https://doi.org/10.3389/ffgc.2021.752063)

Published: 30/08/2021

Publisher's PDF, also known as Version of record

[Cyswllt i'r cyhoeddiad / Link to publication](#)

Dyfyniad o'r fersiwn a gyhoeddwyd / Citation for published version (APA):

Lang, F., Kruger, J., Kaiser, K., Bol, R., & Loeppmann, S. (2021). Editorial: Changes in Forest Ecosystem Nutrition. *Frontiers in Forests and Global Change*, 4. <https://doi.org/10.3389/ffgc.2021.752063>

Hawliau Cyffredinol / General rights

Copyright and moral rights for the publications made accessible in the public portal are retained by the authors and/or other copyright owners and it is a condition of accessing publications that users recognise and abide by the legal requirements associated with these rights.

- Users may download and print one copy of any publication from the public portal for the purpose of private study or research.
- You may not further distribute the material or use it for any profit-making activity or commercial gain
- You may freely distribute the URL identifying the publication in the public portal ?

Take down policy

If you believe that this document breaches copyright please contact us providing details, and we will remove access to the work immediately and investigate your claim.



Editorial: Changes in Forest Ecosystem Nutrition

Friederike Lang^{1*}, Jaane Krüger¹, Klaus Kaiser², Roland Bol^{3,4} and Sebastian Loeppmann^{5,6}

¹ Chair of Soil Ecology, Faculty of Environment and Natural Resources, University of Freiburg, Freiburg, Germany, ² Soil Science and Soil Protection, Martin Luther University Halle-Wittenberg, Halle, Germany, ³ Research Centre Jülich, Institute of Bio- and Geosciences, Agrosphere (IBG-3), Jülich, Germany, ⁴ School of Natural Sciences, Bangor University, Bangor, United Kingdom, ⁵ Biogeochemistry of Agroecosystems, Georg-August-University, Göttingen, Germany, ⁶ Department of Soil Science, Institute of Plant Nutrition and Soil Science, Christian-Albrechts-University, Kiel, Germany

Keywords: phosphorus, nitrogen, cross-scale nutritional interactions, nutrient use efficiency, disturbance, scales of ecosystem nutrition

Editorial on the Research Topic

Changes in Forest Ecosystem Nutrition

OPEN ACCESS

Edited and reviewed by:

Frank Hagedorn,
Swiss Federal Institute for
Forest, Switzerland

*Correspondence:

Friederike Lang
fritzi.lang@
bodenkunde.uni-freiburg.de

Specialty section:

This article was submitted to
Forest Soils,
a section of the journal
Frontiers in Forests and Global
Change

Received: 02 August 2021

Accepted: 05 August 2021


Published: 30 August 2021

Citation:

Lang F, Krüger J, Kaiser K, Bol R and
Loeppmann S (2021) Editorial:
Changes in Forest Ecosystem
Nutrition.
Front. For. Glob. Change 4:752063.
doi: 10.3389/ffgc.2021.752063

Forests strongly depend on natural nutrient resources since fertilization is not a common forest management practice in most parts of the world. Soils and above- and belowground interactions play a crucial role in regulating the retention, distribution, and uptake of nutrients. The high relevance of nutrition for health and productivity of forests has been demonstrated by recent research data obtained by extensive forest monitoring around the world. For example, nutrient availability was the most decisive factor explaining net forest ecosystem productivity (NEP) for a global population of 92 forest sites (Fernández-Martínez et al., 2014). Also, changes in phosphorus (P) and nitrogen (N) nutrition have been emphasized in recent studies on European forests (Jonard et al., 2015; Etzold et al., 2020). These findings stress the necessity of addressing forest nutrition via holistic ecosystem approaches (Figure 1), as was recently outlined for P (Lang et al., 2016).

Linking trees' nutrient supply with the abundance and activity of other organisms present in different ecosystem compartments has a long tradition in forest-related research. Thus, vast information on relationships between soils and site conditions, nutrient cycling and composition, and productivity of vegetation has been collected systematically over long periods of time, and then used to develop empirical models of forest growth (e.g., Barnes et al., 1982). However, forest ecosystem interactions are exposed to quickly changing environmental conditions. Recent drivers of forest nutrition include (a) higher global atmospheric N deposition leading to increased productivity (Etzold et al., 2020) and causing nutrient imbalances (Peñuelas et al., 2012); (b) climate change affecting the recycling of nutrients (Medlyn et al., 2011) and increasing forest disturbance (Gleason et al., 2010); or (c) changes in management intensity leading to increased removal of nutrients with biomass (Vadeboncoeur et al., 2014). Such changes in nutritional conditions are expected to translate into detrimental effects on forest ecosystems. Therefore, proper management of forests for preserving their functioning requires timely updating of empirical knowledge about ecosystem interactions and their feedbacks to nutrition.



Many new approaches of ecosystem nutrition have been compiled in the Research Topic “Changes in Forest Ecosystem Nutrition,” covering spatial scales (Figure 1) from the molecular level (Ganta et al.) up to landscapes (Houle et al.). Those approaches are complementary to the traditional concepts of plant nutrition, which focus on one-way relationships between soil and plants. Key factors in P ecosystem nutrition at discrete scale levels were P speciation in soils (Klotzbücher et al.; Konrad et al.), nutrient mobilization by root exudates (Schack-Kirchner et al.), microbial (Michas et al.), and fungal communities (Müller et al.), identification of P sources and uptake processes (Pistocchi et al.; Hauenstein et al.), P-efficiency of mycorrhizal fungi (Clausing and Polle), the impact of beech provenances (Meller et al.) and root surface distribution (Loew et al.). The topic also addresses factors effective along different scale levels, such as P leaching within soils (Makowski et al.; Gerhard et al.;

Fetzer et al.) or at catchment scale (Sohrt et al.), element coupling during microbial non-steady state conditions at profile scale (Loepmann et al.) or stand scale (Palmqvist et al.), and P heterogeneity at stand scale (Prietzel) and at the scale of landscapes (Yu et al.). In addition, the topic extends to the impact of certain drivers of ecosystem nutrition, including land-use (Heron et al.), pedogenetic and ecosystem succession (Nelson et al.), climate change (Houle et al.) and of N deposition (Braun et al.; Frey et al.).

Overall, the range of publications in the Research Topic impressively illustrates that plant nutrition approaches considering only processes immediately relevant to vitality and yield of individual plants are not sufficient to address recent challenges linked to human impact on forest ecosystems. The importance of saprophytic symbiotic microorganisms – including their community composition, abundance, and activity

– for nutrient turnover in forest ecosystems, and the linkages between above- and below-ground biota, as well as relations to ecosystem services need to be carefully considered. At the scale of ecosystems, heterogeneity and its structuring effects, seasonality, and links to nutrient transport by soil water or within organisms, become crucial factors. Based on the compiled studies, we postulate that for understanding their resilience against and vulnerability to global change, ecosystem nutrition processes need to be characterized based on their emergent interacting, hierarchy, spatial and temporal structure, and robustness. Sound evaluation of the human impact on forest ecosystem nutrition needs assessment beyond nutrient budgets and contents, by considering all relevant key factors that link mobilization and immobilization processes, and therefore shape the nutrient use efficiency of forests at the ecosystem level.

REFERENCES

- Barnes, B. V., Pregitzer, K. S., Spies, T. A., and Spooner, V. H. (1982). Ecological forest site classification. *J. Forestry* 80, 493–498.
- Etzold, S., Ferretti, M., Reinds, G. J., Solberg, S., Gessler, A., Waldner, P., et al. (2020). Nitrogen deposition is the most important environmental driver of growth of pure, even-aged and managed European forests. *Forest Ecol. Manage.* 458:117762. doi: 10.1016/j.foreco.2019.117762
- Fernández-Martínez, M., Vicca, S., Janssens, I. A., Sardans, J., Luysaert, S., Campioli, M., et al. (2014). Nutrient availability as the key regulator of global forest carbon balance. *Nature Climate Change* 4:471–476. doi: 10.1038/nclimate2177
- Gleason, S. M., Read, J., Ares, A., and Metcalfe, D. J. (2010). Species-soil associations, disturbance, and nutrient cycling in an Australian tropical rainforest. *Oecologia* 162, 1047–1058. doi: 10.1007/s00442-009-1527-2
- Jonard, M., Fürst, A., Verstraeten, A., Thimonier, A., Timmermann, V., Potočić, N., et al. (2015). Tree mineral nutrition is deteriorating in Europe. *Global Change Biol.* 21, 418–430. doi: 10.1111/gcb.12657
- Lang, F., Bauhus, J., Frossard, E., George, E., Kaiser, K., Kaupenjohann, M., et al. (2016). Phosphorus in forest ecosystems: New insights from an ecosystem nutrition perspective. *J. Plant Nutr. Soil Sci.* 179, 129–135. doi: 10.1002/jpln.201500541
- Medlyn, B. E., Duursma, R. A., and Zeppel, M. J. B. (2011). Forest productivity under climate change: a checklist for evaluating model studies. *WIREs Clim. Change* 2, 332–355. doi: 10.1002/wcc.108

AUTHOR CONTRIBUTIONS

FL prepared the draft. JK compiled the figures. RB, SL, and KK revised the draft. All authors approved the final submission.

ACKNOWLEDGMENTS

The idea for this Research Topic resulted from the discussion within the Priority Programme 1685 *Ecosystem Nutrition—Forest strategies for limited phosphorus resources* funded by the German Research Foundation (DFG). We greatly thank all authors and reviewers for their contributions to this special issue as well as the support of the editorial office.

Peñuelas, J., Sardans, J., Rivas-ubach, A., and Janssens, I. A. (2012). The human-induced imbalance between C, N and P in Earth's life system. *Global Change Biol.* 18, 3–6. doi: 10.1111/j.1365-2486.2011.02568.x

Vadeboncoeur, M. A., Hamburg, S. P., Yanai, R. D., and Blum, J. D. (2014). Rates of sustainable forest harvest depend on rotation length and weathering of soil minerals. *Forest Ecol. Manage.* 318, 194–205. doi: 10.1016/j.foreco.2014.01.012

Conflict of Interest: The authors declare that the research was conducted in the absence of any commercial or financial relationships that could be construed as a potential conflict of interest.

Publisher's Note: All claims expressed in this article are solely those of the authors and do not necessarily represent those of their affiliated organizations, or those of the publisher, the editors and the reviewers. Any product that may be evaluated in this article, or claim that may be made by its manufacturer, is not guaranteed or endorsed by the publisher.

Copyright © 2021 Lang, Krüger, Kaiser, Bol and Loeppmann. This is an open-access article distributed under the terms of the Creative Commons Attribution License (CC BY). The use, distribution or reproduction in other forums is permitted, provided the original author(s) and the copyright owner(s) are credited and that the original publication in this journal is cited, in accordance with accepted academic practice. No use, distribution or reproduction is permitted which does not comply with these terms.